

**1st and
only**Self-administered Ig
for CIDP maintenance

My Life, My Way With Hizentra

Hizentra is an Ig* therapy that provides
proven CIDP relapse prevention with
the convenience of self-administration,
so you can focus on everyday living

Hizentra is a prescription medicine used to treat CIDP in adults.

*Ig = Immunoglobulin

Please see full Important Safety Information on pages
14–15 and full prescribing information for Hizentra, including
boxed warning and patient product information, in pocket.

Hizentra®
Immune Globulin Subcutaneous
(Human) 20% Liquid

What is CIDP?

Chronic inflammatory demyelinating polyneuropathy (CIDP) is a rare disorder of the nervous system.

- Your immune system generally keeps you healthy by fighting off germs. When you have CIDP, your immune system attacks part of your nerves—a protective layer called myelin
- When myelin is damaged, messages to and from your brain are disrupted and may not get to the rest of your body

Over time this can cause fatigue, a loss of reflexes, or weakness, numbness, or tingling in your arms and legs. Left untreated, CIDP can cause permanent damage to your nerves.

How does Ig therapy help?

Ig therapy is a type of treatment that is infused into the body to help block the immune system from attacking the nerve myelin, though exactly how it works is not completely understood.

- Ig therapy is made from the blood plasma of carefully screened, healthy donors
- While the risk of transmitting infectious agents cannot be completely eliminated, advanced manufacturing processes, including virus reduction steps, are always used in the production of Hizentra

What options are there for infusing Ig treatments?

- 1 Subcutaneous Ig (SCIg) therapy** lets you infuse under your skin, and not into a vein, using a relatively small needle and an infusion pump. Your doctor will provide training to make sure you're prepared to infuse properly, at which point you can infuse at home or wherever your doctor approves.
- 2 Intravenous Ig (IVIg) therapy** is administered by a medical professional into a vein, usually in a hospital or infusion clinic.

Important Safety Information

WARNING: Thrombosis (blood clots) can occur with immune globulin products, including Hizentra. Risk factors can include: advanced age, prolonged immobilization, a history of blood clotting or hyperviscosity (blood thickness), use of estrogens, installed vascular catheters, and cardiovascular risk factors.

If you are at high risk of blood clots, your doctor will prescribe Hizentra at the minimum dose and infusion rate practicable and will monitor for signs of clotting events and hyperviscosity. Always drink sufficient fluids before infusing Hizentra.

See your doctor for a full explanation, and the full prescribing information for complete boxed warning.



What is Hizentra?

Your doctor most likely prescribed an intravenous immunoglobulin (IVIg) therapy to treat your CIDP. This means your treatment is administered with an IV into one of your veins.

Hizentra is an option for adult patients stabilized on IVIg. **It's an immunoglobulin just like IVIg, except it's infused under your skin, not into a vein.** This means you can self-infuse when and where you want, in consultation with your healthcare provider.



"I can live more independently"

– James, on self-administering Hizentra

You and your doctor can choose your dosing schedule based on:



Your routine



How long infusions take



How much medicine per infusion



How many infusion sites

Important Safety Information

Immediately report to your physician any of the following symptoms, which could be signs of serious adverse reactions to Hizentra: Reduced urination, sudden weight gain, or swelling in your legs (possible signs of a kidney problem). Pain and/or swelling or discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, or numbness/weakness on one side of the body (possible signs of a blood clot). Bad headache with nausea; vomiting; stiff neck; fever; and sensitivity to light (possible signs of meningitis). Brown or red urine; rapid heart rate; yellowing of the skin or eyes; chest pains or breathing trouble; fever over 100°F (possible symptoms of other conditions that require prompt treatment).

How is Hizentra different from my IVIg?

Convenient dosing

Hizentra infusions usually take about 1 hour per week.

Infuse where you choose

Self-administration with Hizentra means you won't have to adjust or cancel your plans due to infusion appointments.

Low volume

At 20% concentration, Hizentra is twice as concentrated as 10% IVIg. This means you can infuse a smaller volume per dose.

Easy to use

Most people in the clinical study found Hizentra easy to use.

Important Safety Information


Treatment with Hizentra might not be possible if your doctor determines you have hyperprolinemia (too much proline in the blood), or are IgA-deficient with antibodies to IgA and a history of hypersensitivity. Tell your doctor if you have previously had a severe allergic reaction (including anaphylaxis) to the administration of human immune globulin. Tell your doctor right away or go to the emergency room if you have hives, trouble breathing, wheezing, dizziness, or fainting. These could be signs of a bad allergic reaction.

Does Hizentra work?

Hizentra is proven to prevent CIDP relapse and maintain functional ability

People taking Hizentra:

- Withdrew from the study or relapsed less often than those taking placebo
- Maintained their grip strength, upper-body strength, and lower-body strength



Important Safety Information

Inform your doctor of any medications you are taking, as well as any medical conditions you may have had, especially if you have a history of diseases related to the heart or blood vessels, or have been immobile for some time. Inform your physician if you are pregnant or nursing, or plan to become pregnant.

Is Hizentra safe?

Hizentra demonstrated fewer systemic side effects than IVIg

In clinical trials, Hizentra had fewer systemic side effects per infusion than IVIg. Systemic side effects are those that happen anywhere inside your body, like headache or nausea.

The most common side effects in the clinical trials for Hizentra included redness, swelling, itching, and/or bruising at the infusion site; headache; chest, joint or back pain; diarrhea; tiredness; cough; rash; itching; fever, nausea, and vomiting.

One serious side effect was also reported: allergic dermatitis (skin rash due to allergies).

93%

of Hizentra infusions were completed
without any side effects

There are risks common to all Ig therapies

All Ig treatments, including Hizentra, come with inherent risks, the most serious being thrombosis (blood clotting). **Talk to your doctor to see if you are at risk for thrombosis, and read the Important Safety Information (on pages 14–15 of this brochure) before taking Hizentra.**

Important Safety Information

Infuse Hizentra under your skin *only*; do not inject into a blood vessel. Self-administer Hizentra only after having been taught to do so by your doctor or other healthcare professional, and having received dosing instructions for treating your condition.

How do I infuse Hizentra?


Overview of how to self-administer

A healthcare professional will teach you how to self-infuse Hizentra, and you can view an instructional video or download step-by-step instructions at Hizentra.com.

READY	Wash work surface area and hands before gathering supplies
SET	Prepare Hizentra dose and infusion site(s)
GO	Insert needle and infuse

Hizentra is infused subcutaneously into the fatty tissue just below the skin's first layer. Hizentra should never be infused into a vein.


SCIg needles are relatively small. Depending on your size and weight, your doctor will recommend a needle as short as 1/8" (4 mm) or as long as 1/2" (14 mm).



Important Safety Information


Hizentra is made from human blood. The risk of transmission of infectious agents, including viruses and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent and its variant (vCJD), cannot be completely eliminated.

Needle Sizes



Infusion Sites

- Recommended infusion sites include the stomach, thighs, upper arms, and hips
- Use up to 8 infusion sites simultaneously
 - In clinical trials, most patients used 4 infusion sites
- Infusion sites should be at least 2 inches apart; new sites should be at least 1 inch from a previous site
- Never infuse into areas where the skin is tender, bruised, red, or hard. Avoid infusing into scars or stretch marks



Important Safety Information

The most common side effects in the clinical trials for Hizentra include redness, swelling, itching, and/or bruising at the infusion site; headache; chest, joint or back pain; diarrhea; tiredness; cough; rash; itching; fever, nausea, and vomiting. These are not the only side effects possible. Tell your doctor about any side effect that bothers you or does not go away.

How do I start Hizentra?

If you don't yet have a prescription for Hizentra, you should first talk to your doctor to make sure that Hizentra is right for you. Once you have taken that first important step:

Figure out your treatment details: You and your doctor should work together to decide what dosing schedule works best for you, and where you will get your medicine and supplies. You should receive your supplies from either your specialty pharmacy or your doctor.

Get your Hizentra Welcome Kit: This kit provides important resources to help you start and stay on track with Hizentra. Request one by calling IgIQ or at Hizentra.com.

Learn how to self-infuse: Your healthcare professional will teach you how to infuse Hizentra during your first few weeks of therapy. Be sure to visit Hizentra.com to view a self-administration video or download step-by-step instructions.

Ask your doctor to sign up for Premier StartSM: When your doctor signs you up for the Premier Start sample program, you can receive:

- A free 1-month supply of Hizentra
- Free infusion supplies like tubing, needles, and a pump*
- Free at-home visits by a trained nurse who will teach you how to infuse Hizentra



CSL Behring
premier.startSM



Your single source
for Ig solutions

Call **1-877-355-IGIQ (4447)**
Monday–Friday, 8 AM to 8 PM ET

IgIQ support: Here to help you throughout your journey

IgIQ gathers many Ig resources under one roof, making it easy to get the guidance and **support you need with a single phone call**. Programs include:

Insurance Coverage

IgIQ can help answer questions related to **insurance coverage**. 95% of people with insurance are covered for Hizentra.



The **Hizentra Co-Pay Relief** program helps people who are eligible afford their therapy by helping with monthly out-of-pocket expenses for Hizentra (up to \$5,000 for each year, with no monthly cap[†]).[‡] Most people with commercial insurance pay \$0 out-of-pocket.



CSL Behring Assurance is a program that may help you continue to receive treatment even if you experience a lapse in third-party private health insurance.^{†§}



Voice2Voice is a unique program that connects you with another person with CIDP who can share their personal story of what it's like to live with CIDP and experience Hizentra therapy.^{||}

*Pump must be returned upon completion of sample.

†Other terms and conditions may apply. See program terms and conditions at Hizentra.com.

‡This program applies to product cost only. Contact IgIQ to learn more.

§Does not apply to state- or government-funded programs, such as Medicare or Medicaid.

||Voice2Voice advocates are not healthcare professionals or medical experts. For medical questions, please contact your physician.

Voice2Voice advocates are compensated by CSL Behring LLC for their time and/or expenses.

Before your appointment, ask yourself:

1. How often do you currently receive Ig infusions?

- How many hours does it take to get your Ig infusion? (include travel time if you go to a clinic) _____
- How many days per month do you devote to getting treatment? _____

2. Do you have a port or does your nurse have difficulty finding a vein for IVIg infusions?

3. Have you experienced any IVIg-related side effects? If so, how often? (be sure to report side effects to your physician)

4. Do you ever miss work/school or see decreased productivity due to health problems? If so, how often? _____ days per month (on average)

5. Do you ever feel like you're scheduling your life around your Ig infusions or avoiding making plans altogether?

6. Would you like to have more freedom, control, and flexibility over your Ig administration?

Questions to ask your doctor:

1. What are Ig levels and how would Hizentra affect my Ig levels?

2. What would my treatment look like with Hizentra compared to my current Ig treatment?

- How long would infusions take? What is the best infusion schedule for me?

- Could I travel and still self-infuse? What would I need to consider?

3. Would you recommend Hizentra for me?

For more information about CIDP contact
the GBS | CIDP Foundation at GBS-CIDP.org



Important Safety Information

WARNING: Thrombosis (blood clots) can occur with immune globulin products, including Hizentra. Risk factors can include: advanced age, prolonged immobilization, a history of blood clotting or hyperviscosity (blood thickness), use of estrogens, installed vascular catheters, and cardiovascular risk factors.

If you are at high risk of blood clots, your doctor will prescribe Hizentra at the minimum dose and infusion rate practicable and will monitor for signs of clotting events and hyperviscosity. Always drink sufficient fluids before infusing Hizentra.

See your doctor for a full explanation, and the full prescribing information for complete boxed warning.

Hizentra is a prescription medicine used to treat:

- Primary immune deficiency (PI) in patients 2 years and older
- Chronic inflammatory demyelinating polyneuropathy (CIDP) in adults

Treatment with Hizentra might not be possible if your doctor determines you have hyperprolinemia (too much proline in the blood), or are IgA-deficient with antibodies to IgA and a history of hypersensitivity. Tell your doctor if you have previously had a severe allergic reaction (including anaphylaxis) to the administration of human immune globulin. Tell your doctor right away or go to the emergency room if you have hives, trouble breathing, wheezing, dizziness, or fainting. These could be signs of a bad allergic reaction.

Inform your doctor of any medications you are taking, as well as any medical conditions you may have had, especially if you have a history of diseases related to the heart or blood vessels, or have been immobile for some time. Inform your physician if you are pregnant or nursing, or plan to become pregnant.

Infuse Hizentra under your skin *only*; do not inject into a blood vessel. Self-administer Hizentra only after having been taught to do so by your doctor or other healthcare professional, and having received dosing instructions for treating your condition.

Immediately report to your physician any of the following symptoms, which could be signs of serious adverse reactions to Hizentra:

- Reduced urination, sudden weight gain, or swelling in your legs (possible signs of a kidney problem).
- Pain and/or swelling or discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, or numbness/weakness on one side of the body (possible signs of a blood clot).
- Bad headache with nausea; vomiting; stiff neck; fever; and sensitivity to light (possible signs of meningitis).
- Brown or red urine; rapid heart rate; yellowing of the skin or eyes; chest pains or breathing trouble; fever over 100°F (possible symptoms of other conditions that require prompt treatment).

Hizentra is made from human blood. The risk of transmission of infectious agents, including viruses and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent and its variant (vCJD), cannot be completely eliminated.

The most common side effects in the clinical trials for Hizentra include redness, swelling, itching, and/or bruising at the infusion site; headache; chest, joint or back pain; diarrhea; tiredness; cough; rash; itching; fever, nausea, and vomiting. These are not the only side effects possible. Tell your doctor about any side effect that bothers you or does not go away.

Before receiving any vaccine, tell immunizing physician if you have had recent therapy with Hizentra, as effectiveness of the vaccine could be compromised.

Please see accompanying full prescribing information for Hizentra, including boxed warning and patient product information.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.



Hizentra is the **first** and **only** self-administered immunoglobulin proven to prevent CIDP relapse

Important Safety Information

WARNING: Thrombosis (blood clots) can occur with immune globulin products, including Hizentra. Risk factors can include: advanced age, prolonged immobilization, a history of blood clotting or hyperviscosity (blood thickness), use of estrogens, installed vascular catheters, and cardiovascular risk factors.

If you are at high risk of blood clots, your doctor will prescribe Hizentra at the minimum dose and infusion rate practicable and will monitor for signs of clotting events and hyperviscosity. Always drink sufficient fluids before infusing Hizentra.

See your doctor for a full explanation, and the full prescribing information for complete boxed warning.

Hizentra is manufactured by CSL Behring AG and distributed by CSL Behring LLC.
Hizentra® is a registered trademark of CSL Behring AG.
Biotherapies for Life® and IglQ® are registered trademarks of CSL Behring LLC.
CSL Behring Assurance™ and Premier Start™ are service marks of CSL Behring LLC.
Voice2Voice® is a registered trademark of CSL Behring GmbH.
©2018 CSL Behring LLC 1020 First Avenue, PO Box 61501, King of Prussia, PA 19406-0901 USA
www.CSLBehring.com www.Hizentra.com HIZ-0671-SEP18

Hizentra®
Immune Globulin Subcutaneous
(Human) **20% Liquid**